

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: 83401861.6

(22) Date of filing: 23.09.83

(51) Int. Cl.<sup>3</sup>: C 07 C 69/757

C 07 C 62/20, C 07 C 149/30  
 C 07 C 149/28, C 07 D 333/24  
 C 07 C 91/26, C 07 C 87/30  
 C 07 C 67/343, A 01 N 37/10

(30) Priority: 22.04.83 JP 71264/83

(43) Date of publication of application:  
 31.10.84 Bulletin 84/44

(84) Designated Contracting States:  
 CH DE FR GB IT LI

(71) Applicant: KUMIAI CHEMICAL INDUSTRY CO., LTD.  
 No. 4-26, Ikenohata 1-chome Taio-ku  
 Tokyo(JP)

(71) Applicant: IHARA CHEMICAL INDUSTRY CO., LTD.  
 4-26, Ikenohata 1-chome  
 Taitoh-ku, Tokyo(JP)

(72) Inventor: Motojima, Kenji  
 13-13, Katsuragaoka 3-chome  
 Kakegawa-shi Shizuoka-ken(JP)

(72) Inventor: Miyazawa, Takeshige  
 1809, Kamo Kikugawa-cho  
 Ogasa-gun Shizuoka-ken(JP)

(72) Inventor: Toyokawa, Yasufumi  
 1809, Kamo Kikugawa-cho  
 Ogasa-gun Shizuoka-ken(JP)

(72) Inventor: Matsuzawa, Masafumi  
 132, Ojima Fukude-cho  
 Iwata-gun Shizuoka-ken(JP)


(72) Inventor: Hokari, Hiroshi  
 132, Ojima Fukude-cho  
 Iwata-gun Shizuoka-ken(JP)

(72) Inventor: Kusano, Shoji  
 132, Ojima Fukude-cho  
 Iwata-gun Shizuoka-ken(JP)

(74) Representative: Nony, Michel \*  
 Cabinet Nony 29, rue Cambacérés  
 F-75008 Paris(FR)

(54) New cyclohexane derivatives having plant-growth regulating activities, and uses of these derivatives.

(57) As new compound is provided a cyclohexane derivative  
 of the formula



(I)

wherein R represents a hydrogen atom or an alkyl group, an alkylthioalkyl group or an unsubstituted or substituted phenyl group; and R¹ represents an alkyl group, an unsubstituted or substituted benzyl group, a phenethyl group, a phenoxy-methyl group, a 2-thienylmethyl group, an alkoxymethyl group or an alkylthiomethyl group, or a salt of said cyclohexane compound.

This cyclohexane compound exhibits useful plant-growth regulating effects on crop-plants and also non-crop plants such as lawn and may be prepared by cyclization of an acetonylsuccinic acid dialkyl ester, followed by reaction with an organic acid chloride and by intermolecular rearrangement.

0123001

NEW CYCLOHEXANE DERIVATIVES HAVING PLANT-GROWTH REGULATING  
ACTIVITIES, AND USES OF THESE DERIVATIVES.

Field of the Invention

This invention relates to a new cyclohexane derivative,  
more particularly new derivative of 3,5-dioxo-cyclohexane-  
carboxylic acid, having high plant-growth regulating activities  
5 and to process for the preparation of said cyclohexane  
derivatives. This invention also relates to a plant-growth  
regulating composition comprising said cyclohexane derivative  
as the active ingredient, and further to a method of  
regulating the growth of a plant using said cyclohexane  
10 derivative.


Background of the Invention

Various chemical compounds have been used for regula-  
tion of the growth of plants with a chemical substance, so  
called for the purpose of "chemical control" of the plant-  
15 growth. For example, maleic hydrazide (MH), N,N-dimethyl-  
aminosuccinic amide (known as Daminozide or B-Nine) and  
2-chloroethyl trimethyl ammonium chloride (known as  
Chlormequat or CCC) have been employed for retardation of  
plant-growth, for control of emergence of sideshooting or  
20 for prevention of lodging of plants. These known compounds,  
however, have various drawbacks that their uses are  
restricted in respect of the locus, plants and time which  
can effectively be treated with them, that their plant-  
growth regulating effects are insufficient or unstable,  
25 that they are phytotoxic, and/or that they are too  
expensive.

We have made extensive studies in an attempt to provide new plant-growth regulators which are free from the aforementioned drawbacks of the known plant-growth regulators. As a result, we have now found that certain new cyclohexane derivatives which we have now newly synthesized exhibit some remarkable plant-growth regulating effects and are free from the drawbacks as described above.

# Detailed Description of the Invention

According to an aspect of this invention, therefore, there is provided a cyclohexane compound of the general formula:



wherein R represents a hydrogen atom or an alkyl group, an alkylthioalkyl group or an unsubstituted or substituted phenyl group; and R<sup>1</sup> represents an alkyl group, an unsubstituted or substituted benzyl group, a phenethyl group, a phenoxyethyl group, a 2-thienylethyl group, an alkoxyethyl group or an alkylthioethyl group, or a salt of said cyclohexane compound.


The alkyl group which R may represent in the general formula (I) includes those containing 1 to 8 carbon atoms, for example, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, n-pentyl, and iso-pentyl groups.

The alkyl group which  $R^1$  may represent includes those of 1 to 8 carbon atoms, such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, n-pentyl, n-hexyl, n-heptyl and n-octyl groups.

5 The substituent on the substituted phenyl group which R may represent, as well as the substituent on the substituted benzyl group which  $R^1$  may represent may be a lower alkyl group of 1 to 4 carbon atoms such as methyl; halo group such as chloro; and a lower alkoxyl group of 1 to 4 carbon atoms  
10 such as methoxy.

Typical examples of the cyclohexane compounds of the general formula (I) are listed in Table 1 below, where the physical properties of each compound are also indicated.

Table 1






20








Compound No.	R	$R^1$	Physical Properties
1	$C_2H_5-$	$CH_3-$	m.p. 40 ~ 42°C
2	$n-C_3H_7-$	"	m.p. 43 ~ 44°C
3	$i-C_3H_7-$	"	m.p. 98 ~ 100°C
4	$n-C_4H_9-$	"	$n_D^{20} = 1.4991$
5	$i-C_4H_9-$	"	m.p. 39 ~ 40°C
6	$s-C_4H_9-$	"	m.p. 36 ~ 37°C

to be cont'd...

0123001

5	7	$t\text{-C}_4\text{H}_9\text{-}$	$\text{CH}_3\text{-}$	m.p. 76 ~ 77°C
	8	$\text{CH}_3\text{-}$	$\text{C}_2\text{H}_5\text{-}$	m.p. 49 ~ 50°C
	9	$\text{C}_2\text{H}_5\text{-}$	"	m.p. 55 ~ 56°C
	10	$n\text{-C}_3\text{H}_7\text{-}$	"	$n_D^{20} = 1.4960$
	11	$i\text{-C}_3\text{H}_7\text{-}$	"	m.p. 55 ~ 56°C
10	12	H-	"	m.p. 98 ~ 99°C
	13	"	$n\text{-C}_3\text{H}_7\text{-}$	m.p. 116 ~ 119°C
	14	$\text{CH}_3\text{-}$	"	$n_D^{20} = 1.5025$
	15	$\text{C}_2\text{H}_5\text{-}$	"	m.p. 34 ~ 35°C
	16	$n\text{-C}_3\text{H}_7\text{-}$	"	$n_D^{20} = 1.5010$
15	17	$i\text{-C}_3\text{H}_7\text{-}$	"	$n_D^{20} = 1.4941$
	18	$n\text{-C}_4\text{H}_9\text{-}$	"	$n_D^{20} = 1.4930$
	19	$i\text{-C}_4\text{H}_9\text{-}$	"	$n_D^{20} = 1.4958$
	20	$s\text{-C}_4\text{H}_9\text{-}$	"	$n_D^{20} = 1.4893$
	21	$i\text{-C}_5\text{H}_{11}\text{-}$	"	$n_D^{20} = 1.4885$
20	22		"	m.p. 91 ~ 92°C
	23	$\text{CH}_3\text{-}$ 	"	m.p. 123 ~ 124°C
	24	$\text{Cl-}$ 	"	m.p. 115 ~ 116°C
	25	$\text{C}_2\text{H}_5\text{-}$	$i\text{-C}_3\text{H}_7\text{-}$	$n_D^{20} = 1.4898$
	26	"	$n\text{-C}_4\text{H}_9\text{-}$	$n_D^{20} = 1.4943$
25	27	"	$i\text{-C}_4\text{H}_9\text{-}$	$n_D^{20} = 1.4948$
	28	"	$n\text{-C}_5\text{H}_{11}\text{-}$	$n_D^{20} = 1.4878$
	29	"	$n\text{-C}_6\text{H}_{13}\text{-}$	$n_D^{20} = 1.4803$
	30	"	$n\text{-C}_7\text{H}_{15}\text{-}$	$n_D^{20} = 1.4773$
	31	"	$n\text{-C}_8\text{H}_{17}\text{-}$	$n_D^{20} = 1.4650$

to be cont'd...

5	32	$C_2H_5-$		m.p. 67 ~ 68°C
	33	"		m.p. 80 ~ 81°C
	34	"		m.p. 66 ~ 68°C
	35	"	$C_2H_5OCH_2-$	$n_D^{20} = 1.5035$
	36	"	$C_2H_5SCH_2-$	$n_D^{20} = 1.5357$
	37	H-	$CH_3-$	m.p. 139 ~ 141°C
	38	$CH_3-$	"	m.p. 77 ~ 77.5°C
	39	$t-C_4H_9-$	$C_2H_5-$	m.p. 67 ~ 69°C
	40	$CH_3SC_2H_4-$	$n-C_3H_7-$	$n_D^{20} = 1.5170$
10	41	$C_2H_5-$		m.p. 74 ~ 75°C
	42	"		m.p. 78 ~ 79°C
	43	"		$n_D^{20} = 1.5610$
	44	"		m.p. 61 ~ 63°C


Amongst the particular compounds of Table 1, the following compounds are preferred in this invention.

- (1) Compound No. 8: 3,5-dioxo-4-propionylcyclohexane-carboxylic acid methyl ester;
- (2) Compound No. 9: 3,5-dioxo-4-propionylcyclohexane-carboxylic acid ethyl ester;
- 20 (3) Compound No. 10: 3,5-dioxo-4-propionylcyclohexane-carboxylic acid n-propyl ester;
- (4) Compound No. 13: 3,5-dioxo-4-butyrylcyclohexane-carboxylic acid;
- (5) Compound No. 15: 3,5-dioxo-4-butyrylcyclohexane-carboxylic acid ethyl ester;

- (6) Compound No.16: 3,5-dioxo-4-butyrylcyclohexanecarboxylic acid n-propyl ester;
- (7) Compound No.17: 3,5-dioxo-4-butyrylcyclohexanecarboxylic acid isopropyl ester;
- 5 (8) Compound No.18: 3,5-dioxo-4-butyrylcyclohexanecarboxylic acid n-butyl ester;
- (9) Compound No.21: 3,5-dioxo-4-butyrylcyclohexanecarboxylic acid isopentyl ester;
- (10) Compound No.25: 3,5-dioxo-4-isobutyrylcyclohexanecarboxylic acid ethyl ester; and
- 10 (11) Compound No.37: 3,5-dioxo-4-acetylcyclohexanecarboxylic acid.

For plant-growth regulators to be applied onto grass swards, it is usually required that they should have a particularly improved safety in their stimulus for human's skin and eye since they are frequently used for treatment of swards planted in golf links, parks or yards so that they are likely to be brought into direct contact with human bodies.

20 Through our further studies, we have found that such a class of the new cyclohexane derivative of the general formula (I) which is represented by the following general formula (Ia)




(Ia)

0123001

wherein  $R^2$  represents a hydrogen atom or an alkyl group and  $R^3$  represents an alkyl group can provide such a salt with an inorganic or organic cation which exhibits a significantly reduced stimulus to skin and eye of the animal or human, as compared to the similar salt of the another classes of the new cyclohexane derivative according to the general formula (I), with retaining their excellent plant-growth regulating effects.


According to a preferred embodiment of the first aspect of this invention, therefore, there is provided a salt of a cyclohexane compound of the general formula:



wherein  $R^2$  represents a hydrogen atom or an alkyl group and  $R^3$  represents an alkyl group.

The salt of the new cyclohexane compound according to the general formula (Ia) may be any of the following three types:-

(i) A salt of the cyclohexane compound represented by the general formula:




wherein  $R^2$  and  $R^3$  are as defined above and M represents an



0123001


organic or inorganic cation.

(ii) A salt of the cyclohexane compound represented by the general formula:



5 wherein R<sup>3</sup> and M are as defined above.

(iii) A salt of the cyclohexane compound represented by the general formula:




wherein R<sup>3</sup> and M are as defined above.

10 In the above formulae (Ia), (Ib), (Ic) and (Id), R<sup>2</sup> is as defined hereinbefore and preferably is a hydrogen atom or an alkyl group such as methyl, ethyl, n-propyl, iso-propyl, n-butyl or iso-butyl, R<sup>3</sup> is as defined hereinbefore and preferably is an alkyl group such as mentioned just above,

15 and M stands for an organic or inorganic cation. The inorganic cation here may be the cation of an alkali metal (including ammonium cation), an alkaline earth metal, aluminium, copper, nickel, manganese, cobalt, zinc, iron and silver. The organic cation may be a substituted ammonium


20 ion represented by the formula:



wherein  $R^4$ ,  $R^5$ ,  $R^6$  and  $R^7$  each represents a hydrogen atom or an alkyl, a hydroxyalkyl, an alkenyl, benzyl, a halogen-substituted benzyl, pyridyl or an alkyl-substituted pyridyl group, or  $R^4$  and  $R^5$  taken together form a polymethylene group which may optionally be interrupted by an oxygen atom.

Typical examples of the various salts of the formula (Ib), (Ic) or (Id) according to this invention are shown in Tables 2-4 below.

10 Table 2







(Ib)

15

Compound No.	$R^2$	$R^3$	M	Physical Properties
45	CH <sub>3</sub>	CH <sub>3</sub>	H <sub>2</sub> N <sup>⊕</sup> (C <sub>2</sub> H <sub>4</sub> OH) <sub>2</sub>	m.p. 88 ~ 91°C
46	"	C <sub>2</sub> H <sub>5</sub>	"	m.p. 85 ~ 86°C
47	"	"	H <sub>2</sub> N <sup>⊕</sup> (CH <sub>2</sub> <sup>OH</sup> CHCH <sub>3</sub> ) <sub>2</sub>	m.p. 90 ~ 92°C
48	"	n-C <sub>3</sub> H <sub>7</sub>	H <sub>2</sub> N <sup>⊕</sup> (C <sub>2</sub> H <sub>4</sub> OH) <sub>2</sub>	m.p. 62 ~ 64°C
49	C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	"	(see Table 5 given later)
50	"	"	H <sub>2</sub> N <sup>⊕</sup> (C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub>	(see Table 5)
51	"	C <sub>2</sub> H <sub>5</sub>	"	(see Table 5)
				to be cont'd...



	52	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	H <sub>2</sub> N <sup>⊕</sup> (n-C <sub>3</sub> H <sub>7</sub> ) <sub>2</sub>	m.p. 76 ~ 80.5°C
	53	"	"	H <sub>2</sub> N <sup>⊕</sup> (i-C <sub>3</sub> H <sub>7</sub> ) <sub>2</sub>	m.p. 42 ~ 44.5°C
	54	"	"	H <sub>2</sub> N <sup>⊕</sup> (C <sub>2</sub> H <sub>4</sub> OH) <sub>2</sub>	m.p. 76 ~ 78°C (see Table 5)
	55	"	"	H <sub>2</sub> N <sup>⊕</sup> (CH <sub>2</sub> $\overset{\text{OH}}{\underset{ }{\text{CH}}}$ CH <sub>3</sub> ) <sub>2</sub>	m.p. 91 ~ 94°C
5	56	"	"	H <sub>2</sub> N <sup>⊕</sup> (CH <sub>2</sub> CH=CH <sub>2</sub> ) <sub>2</sub>	(see Table 5)
	57	"	"	H <sub>2</sub> N <sup>⊕</sup> (n-C <sub>4</sub> H <sub>9</sub> ) <sub>2</sub>	m.p. 94 ~ 96°C
	58	"	"	H <sub>2</sub> N <sup>⊕</sup> (i-C <sub>4</sub> H <sub>9</sub> ) <sub>2</sub>	(see Table 5)
	59	"	"	H <sub>2</sub> N <sup>⊕</sup> (CH <sub>2</sub> $\overset{\text{CH}-\text{C}_2\text{H}_5}{\underset{(\text{CH}_2)_3}{\underset{ }{\text{CH}}}}$ ) <sub>2</sub>	refractive index n <sub>D</sub> <sup>25</sup> = 1.4965
	60	"	"	H <sub>2</sub> N <sup>⊕</sup> (n-C <sub>12</sub> H <sub>25</sub> ) <sub>2</sub>	m.p. 62 ~ 63°C
10	61	"	"	n-C <sub>4</sub> H <sub>9</sub> $\searrow$ C <sub>2</sub> H <sub>5</sub> $\nearrow$ N <sup>⊕</sup> H <sub>2</sub>	m.p. 63 ~ 65°C
	62	"	"	H <sub>3</sub> N <sup>⊕</sup> (t-C <sub>4</sub> H <sub>9</sub> )	refractive index n <sub>D</sub> <sup>20</sup> = 1.5069
	63	"	"	HN <sup>⊕</sup> (C <sub>2</sub> H <sub>5</sub> ) <sub>3</sub>	refractive index n <sub>D</sub> <sup>25</sup> = 1.5098
	64	"	"	HN <sup>⊕</sup> (C <sub>2</sub> H <sub>4</sub> OH) <sub>3</sub>	m.p. 53 ~ 56°C
	65	"	"	H <sub>2</sub> <sup>⊕</sup> N 	refractive index n <sub>D</sub> <sup>25</sup> = 1.5505
15	66	"	"	H <sub>2</sub> <sup>⊕</sup> N 	m.p. 55 ~ 56°C
	67	"	"	H <sub>2</sub> <sup>⊕</sup> N 	m.p. 55 ~ 57°C
	68	"	"	H <sub>3</sub> N <sup>⊕</sup> 	m.p. 75 ~ 76°C

0123001



5	69	$C_2H_5$	$C_2H_5$	$H_3N^+$ 	m.p. 67 ~ 68°C
	70	"	"	$1/2Mg^{2+}$	m.p. 184 ~ 188°C
	71	"	"	$1/2Cu^{2+}$	m.p. 203 ~ 206°C (decomp.)
	72	"	"	$1/2Ni^{2+}$	m.p. > 290°C
	73	"	"	$1/2Ca^{2+}$	(see Fig. 1)
10	74	"	"	$Na^+$	(see Fig. 2)
	75	"	"	$K^+$	(see Fig. 3)
	76	"	$n-C_3H_7$	$H_2N^+(C_2H_5)_2$	m.p. 63 ~ 65°C
	77	"	"	$H_2N^+(C_2H_4OH)_2$	m.p. 51 ~ 53°C
	78	"	"	$Na^+$	(see Fig. 4)
15	79	"	"	$K^+$	(see Fig. 5)
	80	"	"	$1/2Ca^{2+}$	m.p. 157 ~ 159°C
	81	"	"	$1/3Fe^{3+}$	refractive index $n_D^{20} = 1.5475$
	82	"	"	$1/3Al^{3+}$	refractive index $n_D^{20} = 1.5294$
	83	$i-C_3H_7$	$C_2H_5$	$H_2N^+(C_2H_4OH)_2$	m.p. 71 ~ 73°C


Table 3



(Ic)


5

10

Compound No.	R <sup>3</sup>	M	Physical Properties
84	n-C <sub>3</sub> H <sub>7</sub>	HN <sup>⊕</sup> (C <sub>2</sub> H <sub>5</sub> ) <sub>3</sub>	refractive index n <sub>D</sub> <sup>20</sup> = 1.5182
85	"	(HOC <sub>2</sub> H <sub>4</sub> )N <sup>⊕</sup> (CH <sub>3</sub> ) <sub>3</sub>	refractive index n <sub>D</sub> <sup>20</sup> = 1.5118
86	"	H <sub>2</sub> N <sup>⊕</sup> (C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub>	refractive index n <sub>D</sub> <sup>20</sup> = 1.5172
87	"	H <sub>2</sub> N <sup>⊕</sup> (C <sub>2</sub> H <sub>4</sub> OH) <sub>2</sub>	refractive index n <sub>D</sub> <sup>20</sup> = 1.5330
88	C <sub>2</sub> H <sub>5</sub>	H <sub>3</sub> N <sup>⊕</sup> (n-C <sub>3</sub> H <sub>7</sub> )	(see Fig. 6)
89	"	(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub> N <sup>⊕</sup> HCH <sub>2</sub> - 	refractive index n <sub>D</sub> <sup>20</sup> = 1.5392
90	"	H <sub>2</sub> N <sup>⊕</sup> (C <sub>2</sub> H <sub>4</sub> OH) <sub>2</sub>	refractive index n <sub>D</sub> <sup>20</sup> = 1.5283
91	"	Na <sup>⊕</sup>	m.p. 225°C (decomp.)
92	"	K <sup>⊕</sup>	m.p. 200 ~ 205°C

0123001

Table 4



(Id)

Compound No.	R <sup>3</sup>	M	Physical Properties
93	C <sub>2</sub> H <sub>5</sub>	Na <sup>⊕</sup>	m.p. 240°C (decomp.)
94	"	H <sub>2</sub> N <sup>⊕</sup> (C <sub>2</sub> H <sub>4</sub> OH) <sub>2</sub>	refractive index n <sub>D</sub> <sup>20</sup> = 1.5351 (see Fig. 7)
95	n-C <sub>3</sub> H <sub>7</sub>	Na <sup>⊕</sup>	refractive index n <sub>D</sub> <sup>20</sup> = 1.5261

In the accompanying drawings, Figs. 1, 2, 3, 4, 5, 6 and 7 show infrared absorption spectra of Compound Nos. 73, 74, 75, 78, 79, 88 and 94, listed in Tables 2-3 respectively.

Table 5 below shows the analytical results of nuclear magnetic resonance spectra (in CDCl<sub>3</sub>) of some of the compounds according to this invention which are indicated with reference to Compound Number as shown in Tables 2-3.

Table 5

Compound No.	Characteristic peaks for particular radicals in N.M.R. Spectra
49	$\begin{array}{c} \text{O} \\   \\ \text{CH}_3\text{CH}_2\text{OC}-: \text{tri, 3H, 1.10 ppm} \end{array}$ $\begin{array}{c} \text{O} \\   \\ \text{CH}_3\text{CH}_2\text{OC}-: \text{q, 2H, 4.13 ppm} \end{array}$ $\begin{array}{c} \text{O} \\    \\ -\text{CCH}_3: \text{s, 3H, 2.33 ppm} \end{array}$ $\text{H}_2\text{N}^{\oplus}(\text{CH}_2\text{CH}_2\text{OH})_2: \text{m, 4H, 3.66 ppm}$ $\text{H}_2\text{N}^{\oplus}(\text{CH}_2\text{CH}_2\text{OH})_2: \text{s, 4H, 4.46 ppm}$ $\left\{ \begin{array}{c} \text{O}^{\ominus}\text{H}_2\text{N}^{\oplus}(\text{CH}_2\text{CH}_2\text{OH})_2 \\   \\ \text{C}_6\text{H}_4 \\   \\ \text{C}=\text{O} \end{array} \right\} : \text{m, 9H, 2.50 - 3.06 ppm}$
50	$\begin{array}{c} \text{O} \\   \\ \text{CH}_3\text{CH}_2\text{OC}-, \text{H}_2\text{N}^{\oplus}(\text{CH}_2\text{CH}_3)_2: \\ \text{tri, 9H, 1.23 ppm} \end{array}$ $\begin{array}{c} \text{O} \\   \\ \text{CH}_3\text{CH}_2\text{OC}-: \text{q, 2H, 4.13 ppm} \end{array}$ $\begin{array}{c} \text{O} \\    \\ -\text{CCH}_3: \text{s, 3H, 2.33 ppm} \end{array}$

to be cont'd...

	$\text{H}_2\text{N}^{\oplus}(\text{CH}_2\text{CH}_3)_2: \text{q}, 4\text{H}, 3.03 \text{ ppm}$ $\text{H}_2\text{N}^{\oplus}(\text{CH}_2\text{CH}_3)_2: \text{s}, 2\text{H}, 9.13 \text{ ppm}$ $\left\{ \text{---} \begin{array}{c} \text{O}^{\ominus} \\ \text{C} \\ \text{C} \\ \text{C} \\ \text{C} \\ \text{C} \end{array} \right. : \text{m}, 5\text{H}, 2.52 \sim 2.82 \text{ ppm}$
51	$\text{CH}_3\text{CH}_2\text{OC}-, -\text{CCH}_2\text{CH}_3, \text{H}_2\text{N}^{\oplus}(\text{CH}_2\text{CH}_3)_2:$ $\text{m}, 12\text{H}, 0.89 \sim 1.39 \text{ ppm}$ $\text{CH}_3\text{CH}_2\text{OC}-: \text{q}, 2\text{H}, 4.15 \text{ ppm}$ $\text{H}_2\text{N}^{\oplus}(\text{CH}_2\text{CH}_3)_2: \text{s}, 2\text{H}, 8.97 \text{ ppm}$ $\left\{ \text{---} \begin{array}{c} \text{O}^{\ominus} \text{H}_2\text{N}^{\oplus}(\text{CH}_2\text{CH}_3)_2 \\ \text{C} \\ \text{C} \\ \text{C} \\ \text{C} \\ \text{C} \end{array} \right. \text{COCH}_2\text{CH}_3: \text{m}, 11\text{H}, 2.52 \sim 3.16 \text{ ppm}$
54	$\text{CH}_3\text{CH}_2\text{OC}-: \text{tri}, 3\text{H}, 1.24 \text{ ppm}$ $\text{CH}_3\text{CH}_2\text{OC}-: \text{q}, 2\text{H}, 4.18 \text{ ppm}$ $\text{H}_2\text{N}^{\oplus}(\text{CH}_2\text{CH}_2\text{OH})_2: \text{m}, 4\text{H}, 3.69 \text{ ppm}$ $\text{H}_2\text{N}^{\oplus}(\text{CH}_2\text{CH}_2\text{OH})_2: \text{s}, 4\text{H}, 6.36 \text{ ppm}$

to be cont'd...



	$\begin{array}{l} \text{O} \\    \\ -\text{CCH}_2\text{CH}_3: \text{tri}, 3\text{H}, 1.09 \text{ ppm} \end{array}$ $\begin{array}{l} \text{O}^\ominus \text{H}_2\text{N}^\oplus (\text{CH}_2\text{CH}_2\text{OH})_2 \\ \{ \text{---} \text{C}_6\text{H}_4 \text{---} \text{COCH}_2\text{CH}_3: \text{m}, 11\text{H}, 2.51 \sim 3.18 \text{ ppm} \end{array}$
56	$\begin{array}{l} \text{O} \\    \\ \text{CH}_3\text{CH}_2\text{OC--}: \text{t}, 3\text{H}, 1.23 \text{ ppm} \end{array}$ $\begin{array}{l} \text{O} \\    \\ \text{CH}_3\text{CH}_2\text{OC--}: \text{q}, 2\text{H}, 4.13 \text{ ppm} \end{array}$ $\begin{array}{l} \text{O} \\    \\ -\text{CCH}_2\text{CH}_3: \text{t}, 3\text{H}, 1.10 \text{ ppm} \end{array}$ $\text{H}_2\text{N}^\oplus -: \text{s}, 2\text{H}, 10.0 \text{ ppm}$ $\text{H}_2\text{N}^\oplus (\text{CH}_2\text{CH}=\text{CH}_2)_2: \text{m}, 6\text{H}, 5.00 \sim 6.16 \text{ ppm}$ $\text{H}_2\text{N}^\oplus (\text{CH}_2\text{CH}=\text{CH}_2)_2: \text{d}, 4\text{H}, 3.26 \text{ ppm}$ $\begin{array}{l} \text{O}^\ominus \\ \{ \text{---} \text{C}_6\text{H}_4 \text{---} \text{COCH}_2\text{CH}_3: \text{m}, 7\text{H}, 2.69 \sim 3.17 \text{ ppm} \end{array}$
58	$\begin{array}{l} \text{O} \\    \\ \text{CH}_3\text{CH}_2\text{OC--}: \text{t}, 3\text{H}, 1.26 \text{ ppm} \end{array}$ $\begin{array}{l} \text{O} \\    \\ \text{CH}_3\text{CH}_2\text{OC--}: \text{q}, 2\text{H}, 4.16 \text{ ppm} \end{array}$

to be cont'd...


5	$\begin{array}{l} \text{O} \\ \parallel \\ -\text{CCH}_2\text{CH}_3: \text{t, } 3\text{H, } 1.08 \text{ ppm} \\ \\ \text{H}_2\text{N}^{\oplus} (\text{CH}_2\text{CH} \begin{array}{c} \text{CH}_3 \\ \text{CH}_3 \end{array})_2: \text{d, } 12\text{H, } 0.93 \text{ ppm} \\ \\ \begin{array}{c} \text{O} \\ \parallel \\ \text{O}^{\ominus} \text{H}_2\text{N}^{\oplus} (\text{CH}_2\text{CH} \begin{array}{c} \text{CH}_3 \\ \text{CH}_3 \end{array})_2 \\   \\ \text{CCH}_2\text{CH}_3: \text{m, } 11\text{H, } 2.52 \sim 3.20 \text{ ppm} \\ \parallel \\ \text{O} \end{array} \\ \\ \text{H}_2\text{N}^{\oplus} -: \text{s, } 2\text{H, } 8.16 \text{ ppm} \\ \\ \text{H}_2\text{N}^{\oplus} (\text{CH}_2\text{CH} \begin{array}{c} \text{CH}_3 \\ \text{CH}_3 \end{array})_2: \text{m, } 2\text{H, } 1.89 \text{ ppm} \end{array}$
---	--

Amongst the particular compounds of Tables 2-4, the following compounds are preferred in this invention.

- (12) Compound No.54: [Bis-(2-hydroxyethyl)]ammonium salt of ethyl 3,5-dioxo-4-propionylcyclohexanecarboxylate.
- (13) Compound No.74: Sodium salt of ethyl 3,5-dioxo-4-propionylcyclohexanecarboxylate.
- (14) Compound No.93: Di-sodium salt of 3,5-dioxo-4-propionylcyclohexanecarboxylic acid.
- (15) Compound No.94: Bis-([bis-(2-hydroxyethyl)]ammonium) salt of 3,5-dioxo-4-propionylcyclohexanecarboxylic acid.

(16) Compound No.55: [Bis-(2-hydroxypropyl)]ammonium salt  
of ethyl 3,5-dioxo-4-propionyl-  
cyclohexanecarboxylate.

The compounds of the general formula (I) can be  
5 prepared by a method briefly depicted according to the  
following reaction equations:-



wherein R and R<sup>1</sup> are as defined hereinabove for the general  
10 formula (I) and M' is an alkali metal.

The 3,5-dioxo-cyclohexanecarboxylic acid alkyl ester  
alkali metal salt of the above formula (III) may be prepared  
by reacting an acetonysuccinic acid di-alkyl ester compound  
of the formula (II) with a basic reagent in a suitable  
15 solvent. This cyclization reaction can be carried out at  
a temperature ranging from room temperature to the boiling  
point of the solvent used for a period of 1 to 10 hours.  
The basic reagent to be used in this reaction includes an  
alkali metal alcoholate such as sodium methylate, sodium

ethylate and potassium tert-butoxide or an alkali metal  
hydride such as sodium hydride. Examples of the solvent  
used here may be methanol, ethanol, benzene, toluene,  
xylene, N,N-dimethylformamide, dimethylsulfoxide and  
5 dimethylcellosolve.

The 3,5-dioxo-cyclohexanecarboxylic acid or its  
alkyl ester compound of the formula (IV) can be made by  
treatment of the compound (III) prepared as above with an  
acid such as hydrochloric acid.


10 The compound of the above formula (VI) can be  
prepared by condensing the compound (III) or (IV) in an  
inert organic solvent with an organic acid chloride of the  
formula (V) where  $R^1$  is as defined hereinbefore, in the  
presence or absence of  $\gamma$ -picoline and in the presence or  
15 absence of a base. This condensation reaction may be carried  
out at a temperature in the range from  $-20^{\circ}\text{C}$  to the boiling  
point of the solvent when used, and preferably at or below  
ambient temperature. The reaction time may vary from 10  
minutes to 7 hours depending upon the reaction conditions  
20 employed. The base which may optionally be used in this  
reaction includes any of such acid-binding agents which are  
generally employed for the conventional dehydrohalogenation,  
for examples, organic bases such as trimethylamine, tri-  
ethylamine, diethylamine, dimethylamine or pyridine and  
25 inorganic bases such as sodium or potassium hydroxide. The  
solvent available for this reaction may include water and/or

organic solvents such as toluene, benzene, xylene, dichloro-  
methane, 1,2-dichloroethane, chloroform, carbon tetrachloride,  
N,N-dimethylformamide, dimethylsulfoxide and methylcellosolve.  
This condensation reaction is conveniently performed with the  
5 aid of  $\gamma$ -picoline which will act as phase transition catalyst  
when using water and organic solvent as the reaction medium.

The new compound of the formula (I) according to this  
invention can be produced by subjecting the intermediate  
compound of the above formula (VI) to a rearrangement  
10 reaction in the presence of a catalyst and an organic  
solvent at a temperature from room temperature to the boiling  
point of the solvent for a period of 1 to 10 hours. The  
catalyst to be used for this rearrangement reaction includes  
pyridine derivatives such as 4-N,N-dimethylaminopyridine,  
15 4-N,N-diethylaminopyridine and 4-pyrolidinoaminopyridine; or  
N-alkylimidazole derivatives such as N-methylimidazole and  
N-ethylimidazole.

The new compound (I) of this invention can be prepared  
without isolation or purification of said intermediate  
20 compounds (III), (IV) and (VI).


According to a second aspect of this invention,  
therefore, there is provided a process for the production  
of a cyclohexane compound of the general formula (I):



(I)


wherein R represents a hydrogen atom or an alkyl group, an alkylthioalkyl group or an unsubstituted or substituted phenyl group; and R<sup>1</sup> represents an alkyl group, an unsubstituted or substituted benzyl group, a phenethyl group, a phenoxyethyl group, a 2-thienylethyl group, an alkoxyethyl group or an alkylthioethyl group, or a salt of said cyclohexane compound, which comprises the steps of:-

(a) reacting a compound of the formula (III):




(III)

or a compound of the formula (IV):




(IV)

wherein R is as defined above and M' is an alkali metal atom, with an organic acid chloride compound of the formula (V):




(V)

wherein R<sup>1</sup> is as defined above, in an organic solvent and/or water as the reaction medium in the presence or absence of a base as an acid-binding agent to produce the compound of the formula (VI):



wherein R and R<sup>1</sup> are as defined above,


- (b) subjecting the compound of the formula (VI) to an intermolecular rearrangement in the presence of a catalyst  
 5 which is chosen from 4-N,N-dimethylaminopyridine, 4-N,N-diethylaminopyridine, 4-pyrrolidinoaminopyridine, N-methylimidazole and N-ethylimidazole to produce the compound of the formula (I):



- 10 wherein R and R<sup>1</sup> are as defined above, and

- (c) if desired, reacting the compound of the formula (I) with an inorganic or organic cation to produce the salt of the compound (I) with said inorganic or organic cation.

As an alternative, such new compounds of the formula  
 15 (I) where R is an alkyl group may be prepared by esterification of the corresponding free carboxylic acid compound according to the following briefed reaction scheme:-



wherein R' represents an alkyl group and  $R^1$  is as defined hereinbefore in the formula (I).

Thus, the compound of the formula (If) can be produced by a method comprising the reaction of the compound of the formula (Ie) with a halogenating agent in the presence or  
5 absence of a solvent and in the presence or absence of a base as an acid-binding agent, followed by a further esterification reaction of the acid halide derivative as formed with an alkanol ( $R'OH$ ) of the formula (VII) in the presence or absence  
10 of a base. The reaction with the halogenating agent may be conducted at a temperature ranging from  $-20^{\circ}C$  to the boiling point of the halogenating agent or of the solvent when used, and preferably at a temperature of from  $-10^{\circ}C$  to  $100^{\circ}C$  for a period of 10 minutes to 7 hours to give an acid halide  
15 derivative of the compound (Ie). The halogenating agent to be used for the above purpose may be thionyl chloride, phosphorus trichloride, phosphorus pentachloride and phosphorus oxychloride. As the solvent which may be used for the halogenation reaction, there may be mentioned  
20 dichloromethane, chloroform, carbon tetrachloride, benzene, toluene, xylene and the like.

The subsequent reaction of esterifying the acid halide derivative with the alkanol (VII) may be carried out at a temperature of  $-20^{\circ}C$  to  $100^{\circ}C$  for a time of 10  
25 minutes to 48 hours. The base which is conveniently used for the halogenation and esterification includes those



which have generally been employed for the conventional dehydrohalogenation as the acid-binding agent in the art and may be selected from organic bases such as triethylamine, pyridine, N,N-dimethylaminopyridine and N,N-dimethylaniline;  
5 and inorganic bases such as sodium hydroxide, potassium carbonate and sodium bicarbonate. The esterification with the alkanol (VII) may be conducted after or without separation of the intermediate acid halide derivative from the reaction medium.


10           The salts of the new cyclohexane compounds of the formula (I) or (Ia) according to this invention can be prepared as follows:-

          Thus, for example, an organic or inorganic salt of the new cyclohexane compound may be produced by reacting in  
15 an organic solvent a compound of the general formula (I) or (Ia) with one or two or more equivalent proportions of a salt-forming reagent which is chosen from a primary, secondary or tertiary amine, a metal alcoholate, or a metal salt such as a chloride, sulfate, nitrate, acetate  
20 or carbonate; or hydride or hydroxide of a metal, for example, an alkali metal such as sodium, potassium; an alkaline earth metal such as calcium, magnesium, barium; aluminium, nickel, copper, manganese, cobalt, zinc, iron or silver. When the compound (I) or (Ia) is reacted with  
25 one equivalent proportion of the salt-forming reagent, either the salt formed will be in the form of the salt of

0123001


the general formula (Ib) as indicated in Table 2 when the group R or R<sup>2</sup> in the compound of the formula (I) or (Ia) is alkyl, or the salt formed will be in the form of the salt of the general formula (Ic) as indicated in Table 3 when R or R<sup>2</sup> is hydrogen in the compound (I) or (Ia). If two or more equivalent proportions of the salt-forming reagent is reacted with the compound (I) or (Ia), the salt as formed will take the form of the general formula (Id) as indicated in Table 4, when R or R<sup>2</sup> is a hydrogen atom in the compound (I) or (Ia).

The new compounds of the formula (I) or (Ia) according to this invention may possibly undergo a tautomerism and may be present in the form of tautomeric isomers as shown below, which fall within the scope of this invention.



wherein Z denotes R or R<sup>2</sup> as defined hereinbefore and Y denotes R<sup>1</sup> or R<sup>3</sup> as defined hereinbefore.


Further, in an aqueous solution, the new compound salt of the formula (Ib) or (Ic) according to this invention may possibly change as shown below:-



5 It may be added that when the cyclohexane compound of the formula (Ia) where  $R^2$  is an alkyl group, is reacted with a primary, secondary or tertiary amine which may be represented by the formula




10 where  $R^4$ ,  $R^5$  and  $R^6$  are each as defined hereinbefore but all of  $R^4$ ,  $R^5$  and  $R^6$  cannot be each a hydrogen atom simultaneously, there is formed such salt of the compound (Ia) which may be deemed as the substituted ammonium salt of the formula



15 Further, when the cyclohexane compound of the formula (Ia) where  $R^2$  is a hydrogen atom is reacted with a primary, secondary or tertiary amine of the formula (VIII), there is

0123001

formed such salt of the compound (Ia) which may also be deemed as the substituted ammonium salt of the formula



In Tables 2-4, the salts of the new compounds with  
5 amine are represented as the type of the substituted ammonium salt as shown just above.

The plant-growth regulating effects of the new compounds of the general formula (I) and the salts thereof are manifested predominantly as a stunting or dwarfing effect on the  
10 vegetative growth of plants, but other various plant-growth regulating effects may be manifested by modifying the nature of plants to be treated and the means, times and rates of application of the compounds or salts. Such plant-growth regulating effects which may be induced by the new compound  
15 of this invention include promotion of rooting, reduction in risk of lodging, promotion of sideshooting and root growth, maintenance of green color of stems and leaves, promotion or delay of flowering, promotion of fruiting and ripening, and increase in resistances to temperature hindrance, to  
20 phytotoxicity caused by herbicides and to fungal or bacterial diseases. These various effects are not always manifested at a time.

The new compounds (including the salts) of this

0123001

invention can control internode elongation of cereals and prevent or reduce lodging of rice, wheat, barley, maize and the other crop-plants by foliage and/or soil-treatment. Application of the compound of this invention to seedlings  
5 of aquatic rice and growing vegetables can lead an improvement in quality, rooting after transplanting, and resistance to low temperature.

The foliage treatment with the new compound during the vegetative stage of crop plants, e.g. aquatic rice,  
10 wheat and barley also induces shortening of top leaves or improvement in leaf orientation and hence increases in light interception and utilization which will enhance the ripening and increase the overall grain yield of such crop-plants.

Moreover, the compound of this invention may suppress  
15 spindly growth of flowers, ornamental plants and horticultural plants which may be caused due to high temperature or sunshine shortage in plantation in a greenhouse.

The compound of the invention which exhibits the above-mentioned plant-growth regulating effects are very  
20 useful not only for agricultural and horticultural treatment but also for control of plant growth in non-crop lands. For instance, when the compound of this invention is applied onto lawns in park, playing field, golf link, airport or embankment or undergrowth grasses in orchard or pasture  
25 land, it is possible to inhibit the overluxuriant growth, to reduce the number of reaping and/or to facilitate the

mowing operations as usually required for maintenance. Further, application of the new compound of this invention onto swards can promote sideshooting and increase the planting density of swards.

5           The compounds of this invention may be used as such for plant-growth regulating purposes but are more conveniently formulated into compositions for such usage. According to a third aspect of this invention, there is thus provided a plant-growth regulating composition comprising  
10           a compound of the general formula (I) as hereinbefore defined or a salt thereof in combination with a carrier or diluent.

          The invention also provides a method of regulating the growth of plant, which comprises applying an effective  
15           amount of a cyclohexane compound of the formula (I) or a salt thereof of this invention to the foliage or seed of the plant to be treated or to the soil or locus where the plant to be treated is grown.

          The compounds and salt can be applied in a number of  
20           ways, for example, they can be applied, formulated or unformulated, directly to the foliage of a plant, or they can be applied also to bushes and trees, to seeds or to other medium in which plants, bushes or trees are growing or are to be planted, or they can be sprayed on, dusted on  
25           or applied as a cream or paste formulation, or they can be applied as slow release granules.

The compounds are preferably used for agricultural and horticultural purposes in the form of a composition. The type of composition used in any instance will depend upon the particular purpose envisaged.

5           The compositions may be in the form of dusting powders or granules comprising the active ingredient and a solid diluent or carrier, for example fillers such as kaolin, bentonite, dolomite, calcium carbonate, talc, powdered  
10           magnesia, Fuller's earth, gypsum, diatomaceous earth and China clay.

          The compositions may also be in the form of dispersible powders, granules or grains comprising a wetting agent to facilitate the dispersion in liquids of the powder or grains which may contain also fillers and suspending  
15           agents.

          The aqueous dispersions or emulsions may be prepared by dissolving the active ingredient(s) in an organic solvent optionally containing wetting, dispersing or emulsifying agent(s) and then adding the mixture to water which may also  
20           contain wetting, dispersing or emulsifying agent(s). Suitable organic solvents are kerosene, cyclohexanone, methylethyl ketone, acetone, methanol and acetonitrile.

          The compositions may also be in the form of liquid preparations for use as dips or sprays which are generally  
25           aqueous dispersions or emulsions containing the active ingredient in the presence of one or more of wetting

agent(s), dispersing agent(s), emulsifying agent(s) or suspending agent(s). These agents can be anionic or non-ionic agents.

5 The compositions for use as aqueous dispersions or emulsions are generally supplied in the form of a concentrate containing a high proportion of the active ingredient(s), and the concentrate is to be diluted with water before use.

10 The plant-growth regulating composition of this invention may usually be formulated into a wettable powder comprising 5 to 95%, preferably 10 to 50% by weight of the new compounds of this invention as active ingredient; 1 to 20%, preferably 5 to 10% by weight of surfactant; and 4 to 44%, preferably 40 to 85% by weight of solid carrier.

15 The composition of this invention may be formulated into an emulsifiable concentrate comprising 5 to 95%, preferably 20 to 70% by weight of the new compound of this invention as active ingredient; 1 to 40%, preferably 5 to 20% by weight of surfactant; and 4 to 94%, preferably 10 to 75% by weight of liquid carrier.

20 The composition of this invention may be made up as granules comprising 0.5 to 40%, preferably 2 to 10% by weight of the new compound of this invention as active ingredient; 1 to 20%, preferably 2 to 10% by weight of surfactant; and 40 to 98.5%, preferably 80 to 96% by weight  
25 of solid carrier. And, the composition of this invention may be formulated into dust comprising 0.5 to 10%,



preferably 1 to 5% by weight of the active ingredient; and 99.5 to 90%, preferably 99 to 95% by weight of finely divided solid carrier.

The composition of this invention may also be formulated into a paste comprising 0.1 to 20%, preferably 1 to 10% by weight of the active ingredient, 1 to 20%, preferably 2 to 10% by weight of surfactant; and 60 to 98.9%, preferably 80 to 97% by weight of paste base.

The rate of application of the new compounds of this invention for use in the purposes of plant-growth regulation will normally depend on various factors, including nature of plants to be treated, application time and application means. In general, however, the rate of application may be in the range of 0.01 kg to 50 kg per hectare and preferably of 0.05 kg to 10 kg per hectare as the active ingredient.

For the purpose of preventing or reducing lodging of crop plants such as wheat, the rate of application may desirably be in a range of 0.1 kg to 1 kg per hectare as the active ingredient. For the purpose of suppressing the growth of swards in non-crop lands or undergrowth-grasses in orchards, the rate of application may be in a range of 0.5 kg to 5 kg per hectare as the active ingredient. For the purpose of regulating the growth of aquatic rice plants, the rate of application may be in the range of 0.5 kg to 5 kg per hectare as the active ingredient when it is applied to the soil, and the formulations to be applied to the

0123001

foliage of aquatic rice plant may contain the active ingredient at a concentration of 100 ppm to 5000 ppm.

In general, the composition of this invention may be formulated into a diluted or dilutable preparation containing the active  
5 ingredient at a concentration of 10 ppm to 10,000 ppm and preferably of 100 ppm to 5000 ppm for use in the foliage treatment.

This invention is now illustrated with reference to the following Examples to which this invention is not  
10 limited.

Example 1

Preparation of Compound No. 15, namely ethyl 3,5-dioxo-  
-4-butyrylcyclohexanecarboxylate.

To 150 ml of ethanol containing 3.4 g of sodium was  
15 added dropwise 31 g of diethyl acetonylsuccinate (or diethyl acetmethylsuccinate) over about one hour with stirring at ambient temperature. The mixture was heated under reflux for 2 hours and then cooled to room temperature, followed by distillation off of the ethanol under reduced pressure.  
20 There was thus obtained a brown viscous liquid comprising sodium salt of ethyl 3,5-dioxocyclohexanecarboxylate as formed, to which were added 100 ml of toluene, 50 ml of ice water and 0.3 g of  $\gamma$ -picoline and then added dropwise 15 g of n-butyryl chloride at ambient temperature under agitation  
25 over about 2 hours. The resultant reaction mixture was agitated for further 30 minutes and then the toluene layer

was separated, washed with water, dried and admixed with 0.8 g of 4-N,N-dimethylaminopyridine. The admixture was heated with agitation at a temperature of 80-90°C for 3 hours. The toluene was then distilled off from the reaction solution under reduced pressure and the residue was chromatographed on a column of silica gel developed with benzene to give 19.5 g (55.9%) of the object compound (Compound No. 15).

Compound Nos. 1, 2 and Nos. 4-44 could be prepared by the same procedure as described in Example 1.

Example 2

Preparation of Compound No. 26, namely ethyl 3,5-dioxo-4-n-valerylcylohexanecarboxylate.

2.1 g of 5-ethoxycarbonyl-cyclohexane-1,3-dione, namely ethyl 3,5-dioxocyclohexanecarboxylate was dissolved in 20 ml of dichloromethane, to which was added 1.7 g of triethylamine and then added dropwise 1.9 g of n-valeryl chloride with stirring over 30 minutes. The mixture was continued to be stirred for 3 hours, and the reaction solution obtained was washed with water. The organic layer was separated from the aqueous phase, dried and concentrated to a small volume.

The concentrated solution was taken up in 20 ml of toluene, followed by addition of 0.6 g of 4-N,N-dimethylaminopyridine and heating at a temperature of 80-90°C for 4 to 5 hours. Subsequently, the toluene was removed from

0123001


the reaction solution by distillation under reduced pressure, and the residue was chromatographed on silica gel using benzene as the developing solvent to yield 2.2 g (71.9%) of the aimed compound (Compound No. 26).

5           Compound Nos. 1-25 and Nos. 27-44 could be produced by the same procedure as stated in Example 2.

Example 3

Preparation of Compound No. 51 (see Table 2)

1.0 g of ethyl 3,5-dioxo-4-propionylcyclohexane-  
10   carboxylate was dissolved in 20 ml of chloroform, to which was then added 1.0 g of diethylamine as the salt-forming reagent. The mixture was allowed to stand at room temperature for two hours, after which the chloroform and the excess of diethylamine were distilled off from the reaction solution  
15   under reduced pressure to yield 1.3 g of the object compound (Compound No. 51) as a pale brown solid and represented by the formula:



Example 4

20           Preparation of Compound No. 54 (see Table 2)

4.0 g of ethyl 3,5-dioxo-4-propionylcyclohexane-  
carboxylate was dissolved in 50 ml of ethanol and 1.7 g of diethanolamine was added to the solution, followed by

stirring for 30 minutes. The ethanol was then distilled off from the reaction solution under reduced pressure to precipitate a crystal, which was washed with ethyl ether and dried to afford 5.2 g of the object compound as a pale yellow  
5 crystal having a melting point of 76-78°C.

Example 5

Preparation of Compound No. 63

1.0 g of triethylamine was added to 1.0 g of ethyl 3,5-dioxo-4-propionylcyclohexanecarboxylate dissolved in  
10 20 ml of chloroform, and the resultant mixture was allowed to stand at ambient temperature for 2 hours. The chloroform and the excess triethylamine were distilled off from the reaction solution under reduced pressure to yield 1.4 g of the desired compound (Compound No. 63) in the form of brown  
15 viscous liquid with a refractive index  $n_D^{20} = 1.5098$ .

Example 6

Preparation of Compound No. 73

1.0 g of potassium salt of ethyl 3,5-dioxo-4-propionylcyclohexanecarboxylate was dissolved in water, to  
20 which was added 0.25 g of calcium chloride. The mixture was allowed to stand overnight to effect the cation-exchange reaction, after which the reaction solution was concentrated by evaporation of the water. The concentrated solution was extracted with hot ethanol and the ethanolic extract was  
25 admixed with hexane to precipitate a crystal. The latter was filtered and dried to give 0.7 g of the aimed compound

as a colorless powder having a melting point of 213-215°C.

Example 7

Preparation of Compound No. 74

To 8.4 g of ethyl 3,5-dioxo-4-propionylcyclohexane-  
5 carboxylate dissolved in 80 ml of ethyl ether was added  
dropwise under agitation at room temperature 50 ml of  
ethanolic solution containing 1.4 g of sodium hydroxide.  
After completion of the addition, the precipitated crystal  
was collected by filtration and dried. Recrystallization  
10 from a mixed solvent of ethanol and hexane gave 7.2 g of  
the desired sodium salt compound as colorless needles with  
a melting point of 178.5-179.5°C.

Example 8

Preparation of Compound No. 78

15 2.0 g of ethyl 4-butyryl-3,5-dioxo-cyclohexane-  
carboxylate was added into ethanol containing 0.53 g of  
sodium ethylate, and the mixture was allowed to stand at  
room temperature for 3 hours. The ethanol was then distilled  
off from the reaction solution under reduced pressure to  
20 yield 2.17 g of the desired sodium salt as a pale yellow  
amorphous solid.

Example 9

Preparation of Compound No. 84 (see Table 3)

1.3 g of 4-butyryl-3,5-dioxo-cyclohexanecarboxylic  
25 acid was dissolved in 20 ml of methanol and 0.58 g of  
triethylamine was added to the solution. The mixture was

0123001


allowed to stand overnight at ambient temperature, after which the methanol was removed from the reaction solution by evaporation under reduced pressure to leave 1.8 g of the object compound (Compound No. 84) as a pale yellow viscous liquid with  $n_D^{20} = 1.5182$ .

The above procedure was repeated using choline in place of the triethylamine to give Compound No. 85.

#### Example 10

##### Preparation of Compound No. 87

1.0 g of 4-butyryl-3,5-dioxo-cyclohexanecarboxylic acid was dissolved in 20 ml of methanol and 0.4 g of diethanolamine  $(HOCH_2CH_2)_2NH$  was added to the solution. The mixture was allowed to stand at room temperature for 3 hours and the methanol was then removed from the reaction solution by evaporation under reduced pressure to give 1.4 g of the desired compound as a pale brown viscous liquid with  $n_D^{20} = 1.5330$ . This product was of the formula:



#### Example 11

##### Preparation of Compound No. 91

0.84 g of sodium bicarbonate was added to 2.1 g of 3,5-dioxo-4-propionylcyclohexanecarboxylic acid dissolved in 30 ml of ethanol, and the mixture was heated under reflux for 3 hours. The reaction solution was then cooled and

filtered to leave a crystal, which was recrystallized from ethanol to give 2.2 g of the desired compound as pale yellow flakes with a melting point of 225°C (decomp.).

Example 12

5     Preparation of Compound No. 93 (see Table 4)

1.4 g (6.6 mmoles) of 3,5-dioxo-4-propionylcyclohexanecarboxylic acid was dissolved in 30 ml of ethanol, and 0.7 g (6.6 mmoles) of anhydrous sodium carbonate was added to the solution. The resultant mixture was heated under reflux for  
10 6 hours, after which the reaction solution was cooled and filtered to leave a crystal. Recrystallization of the crystal from a mixed solvent of ethanol, methanol and hexane gave 1.3 g of the desired di-sodium salt compound as a pale yellow powder having a melting point of 240°C (decomp.).

15     Example 13

Preparation of Compound No. 95

2.0 g of 4-butyryl-3,5-dioxo-cyclohexanecarboxylic acid was dissolved in a mixture of methanol and water (1:1), to which was then added 0.72 g of sodium hydroxide. After  
20 complete dissolution of the latter, the resulting solution was allowed to stand overnight, followed by removal of the methanol and water by distillation under reduced pressure. There was thus obtained 2.4 g of the object compound as a redish brown viscous liquid with  $n_D^{20} = 1.5261$ .



Example 14

Preparation of ethyl 3,5-dioxo-4-propionylcyclohexane-  
carboxylate

4.2 g of ethyl 3,5-dioxocyclohexanecarboxylate and  
5 3.4 g of triethylamine were dissolved in 50 ml of dichloro-  
methane, to which was added dropwise 2.9 g of propionyl  
chloride under stirring at room temperature. The resultant  
mixture was allowed to react at ambient temperature for 2  
hours, after which the reaction solution was washed with  
10 water and admixed with a dilute aqueous sodium bicarbonate  
solution to remove the acidic matters from the reaction  
solution. The organic layer was then separated from the  
aqueous phase, dried and concentrated to a small volume.  
The concentrated solution was taken up in 20 ml of toluene,  
15 followed by addition of 0.5 g of 4-N,N-dimethylaminopyridine  
and heating the mixture at 85°C for 3 hours. Thereafter,  
the toluene was removed from the reaction solution by distil-  
lation under reduced pressure, and the residue was  
chromatographed on silica gel using benzene as developing  
20 solvent to afford 2.5 g of the title compound (Compound  
No. 9 of Table 1) as colorless flakes having a melting point  
of 55-56°C.

Example 15

Preparation of ethyl 4-n-butyryl-3,5-dioxocyclohexane-  
25 carboxylate

28.8 g of sodium salt of ethyl 3,5-dioxocyclohexane-

carboxylate was admixed with 100 ml of toluene and 50 ml of ice water, and to the resulting suspension was added 0.3 g of  $\gamma$ -picoline. To the mixture obtained was added dropwise 15 g of n-butyryl chloride with stirring over about two hours, and the resultant mixture was continued to be stirred for 30 minutes. The toluene layer was then separated from the aqueous phase, dried and concentrated to a small volume. The concentrated solution was dissolved in 20 ml of toluene and 0.8 g of 4-N,N-dimethylaminopyridine was added to the solution, which was then heated at 80-90°C under agitation for 3 hours. Thereafter, the toluene was removed from the reaction solution by distillation under reduced pressure, and the residue was chromatographed on silica gel developing with benzene to yield 19.5 g of the title compound (Compound No. 15 of Table 1) as pale yellow prisms with a melting point of 34-35°C.

The following Examples 16-20 illustrate the preparation of the compositions according to this invention, in which parts are all by weight.

20 Example 16

Wettable powder

A composition in the form of wettable powder was prepared by mixing homogeneously and grinding 10 parts of Compound No. 1, 85 parts of kieselguhr, 2 parts of sodium dinaphthylmethanesulfonate and 3 parts of sodium lignosulfonate.

0123001

A similar wettable powder was produced by the same procedure as above but using Compound No. 45 instead of Compound No. 1.

Example 17

5      Emulsifiable concentrate

An emulsifiable concentrate was made up by mixing the ingredients, and stirring the mixture until all the constituents were dissolved.

	Compound No. 15	30 parts
10	Cyclohexane	20 parts
	Polyoxyethylenealkylaryl ether	11 parts
	Calcium alkylbenzenesulfonate	4 parts
	Methylnaphthalene	35 parts

A similar emulsifiable concentrate was prepared by  
15 the same procedure as described above but replacing Compound No. 15 by Compound No. 46.

Example 18

Grains

A composition in the form of grains was prepared by  
20 uniformly mixing and grinding together 5 parts of Compound No. 13, 2 parts of sodium laurylsulfate, 5 parts of sodium lignosulfonate, 2 parts of carboxymethylcellulose and 86 parts of clay, followed by addition of 20 parts of water per 80 parts of the mixture. The resultant mixture was  
25 milled and formulated into grains of 14-32 mesh size by means of an extrusion granulator.

0123001

A similar granular composition was made up by repeating the above procedure but replacing Compound No. 13 by Compound No. 47.

Example 19

5     Dusting powder

4 parts of Compound No. 16, 5 parts of kieselguhr and 91 parts of clay were homogeneously mixed and ground together to formulate into a dusting powder.

The above procedure was repeated using Compound No. 48  
10 instead of Compound No. 16 to produce a similar dusting powder.

Example 20

Paste formulation

A paste formulation was prepared by mixing uniformly 5 parts of Compound No. 9, 1 part of xylene, 5 parts of  
15 polyoxyethylenesorbitan monolaurate and 89 parts of white vaseline as paste base.

A similar paste formulation was made up by the same procedure as mentioned above except that Compound No. 9 was replaced by Compound No. 49.

20     The following Examples illustrate the plant-growth regulating effects of the compounds according to this invention.

Example 21

This Example illustrates the retardation effect of  
25 the compounds of this invention on the growth of aquatic rice plants.

Aquatic rice plants ("Kinmage", Variety) at the two-  
leaf stage were treated with the compounds of this invention  
under test. Thus, the compounds were formulated into a  
wetttable powder as prepared in Example 16 and the wetttable  
5 powder was diluted with such a volume of water that 1000 g  
of the diluted aqueous preparations as sprayed gave the rate  
of application of 1 kg of the active ingredient per hectare  
upon the foliage treatment by spraying.

On the day of the application and 3 weeks after the  
10 application of the compounds, the length of stems and leaves  
of the treated plants were measured, and retardation of  
growth was calculated by the following equation:-

$$\text{Retardation (\%)} = \left(1 - \frac{B}{A}\right) \times 100$$

A = (Foliar length in untreated plots 3 weeks after  
15 application) - (Foliar length in untreated plots  
on the day of application)

B = (Foliar length in treated plots 3 weeks after  
application) - (Foliar length in treated plots on  
the day of application)

20 The test results are set out in Table 6 below.

Table 6

	Compound No. tested	Retardation (%)	Compound No. tested	Retardation (%)
	1	57	24	63
	2	58	25	81
5	3	57	26	55
	4	64	27	56
	5	67	28	49
	6	52	29	41
	7	49	30	30
10	8	67	31	22
	9	71	32	61
	10	78	33	63
	11	65	34	58
	12	74	35	57
15	13	68	36	76
	14	59	37	74
	15	82	38	62
	16	79	39	61
	17	56	40	53
20	18	58	41	54
	19	66	42	58
	20	69	43	57
	21	77	44	61
	22	63	45	53
25	23	62	46	63

to be cont'd...



0123001

5	47	65	73	66
	48	52	74	72
	49	46	75	70
	50	44	76	62
	51	61	77	65
10	52	60	78	59
	53	62	79	63
	54	64	80	57
	55	63	81	72
	56	60	82	69
15	57	62	83	66
	58	59	84	70
	59	65	85	49
	60	64	86	58
	61	65	87	58
20	62	60	88	61
	63	57	89	68
	64	60	90	57
	65	65	91	73
	66	63	92	70
25	67	62	93	65
	68	70	94	71
	69	66	95	67
	70	67	CCC (Comparative)	0
	71	64	B-Nine (Comparative)	0
	72	71	MH (Comparative)	41

The results of Table 6 indicate that rice plant height can be controlled by the application of the new compounds of this invention and thus spindly growth of rice plant can be prevented.

5 Example 22

This Example illustrates the stunting effect of the compounds of this invention on the growth of grass swards.

Creeping bentgrass (Agrostis stolonifera) ("Seaside" Variety) and Japanese lowngress (Zoysia japonica) of 2 cm  
10 in height were sprayed with the diluted aqueous preparations containing the compounds of this invention. The diluted aqueous preparations as used were prepared by formulating the compounds into wettable powder as in Example 16 and  
15 diluting the wettable powder with such a volume of water that 1000 l of the diluted aqueous preparations as sprayed gave the rate of application of 1 kg of the active ingredient per hectare upon the foliage treatment by spraying.

5 Weeks after application of the compounds, assessment was made and percentage of retranslocation was calculated in the  
20 same way as described in Example 21. Phytotoxicity was also assessed for some of the test compounds and indicated by the following scales:

	<u>Scales</u>	<u>Degree of Phytotoxicity</u>
	-	Not detected
25	±	Small
	+	Middle
	++	Great
	+++	Kill



0123001

The test results are shown in Tables 7a and 7b below.

Table 7a

Compound No. tested	Retardation (%)	
	Creeping bentgrass	Japanese lowgrass
1	-	38
9	65	65
13	70	71
15	62	63
16	-	42
17	-	40
18	-	54
25	60	75

Table 7b

Compound No. tested	Retardation (%)		Phytotoxicity	
	Creeping bentgrass	Japanese lowgrass	Creeping bentgrass	Japanese lowgrass
46	45	68	-	-
54	51	58	-	-
74	53	59	-	-
90	49	54	-	-
94	40	43	-	-
MH (Com- parative)	-	37	++	+

From the results of Tables 7a and 7b as above, it will be seen that the new compounds and salts of this invention exert the stunting or dwarfing effects on lawn grasses for a long period without significant phytotoxic effects and without deteriously affecting the appearance (particularly the color) of the grass, so that the compounds of this invention are useful to reduce the need for mowing the grass swards.

Example 23

10           This Example illustrates the improvements in quality of rice seedlings by application of the compounds of this invention.

Emergent seeds of aquatic rice ("Kinmaze" Variety) were sown in a rice-planting case of 30 x 60 cm in size, then covered with soil and irrigated with water. The compounds under test were formulated into an emulsifiable concentrate as stated in Example 17 which was then diluted with water so that the diluted emulsions contained the active ingredient at the concentrations as indicated in Table 8 below. The diluted emulsions so formulated were sprayed onto the soil surface at the rate of 20 ml per case. The cases were then kept in a thermostatic chamber at 30°C for 3 days and further placed in a greenhouse. When the rice seedlings in untreated plots had reached the stage of 2.5 leaf age, the height of seedlings, leaf age and dry weights of the foliage parts (stems/leaves) and roots of the treated

15

20

25

and untreated plants were measured. Some members of the seedlings were washed with water, and then the seedlings were transplanted into and placed at an angle of 45° in a pot flooded with water to 2 cm depth, after cutting the  
5 roots into the constant length of 2 cm. 10 Days later, the rising angle of the seedlings and the chlorophyll content of their second leaf were determined.

The results are shown in Table 8 below, where the assessment particulars other than the rising angle are  
10 indicated in terms of the percentages based on the results obtained for the untreated plots.

It will be seen from Table 8 that the compounds of this invention can suppress the excessive growth of rice seedlings in a planting case, promote the growth of roots  
15 and accelerate the rooting of transplanted rice seedlings.

Table 8

Compound No. tested	Concentration (ppm) of active ingredient in the sprayed emulsion	Quality of rice seedling at the time transplantation				Quality of rice seed- ling 10 days after transplantation	
		Plant height (%)	Leaf age (%)	Dry weight of foliage parts (%)	Dry weight of root parts (%)	Raising angle of seedling (degree)	Chlorophyll content of second leaf (%)
15	4000	59	100	92	113	90	111
	2000	67	100	97	119	90	110
	1000	88	100	101	120	87	108
54	1000	83	104	98	110	90	113
55	1000	80	108	96	115	90	120
57	1000	85	104	98	112	90	112
70	1000	79	108	96	119	90	136
87	1000	76	108	94	119	90	139
93	1000	83	104	97	112	90	118
Untreated	-	100	100	100	100	75	100

Example 24

This Example illustrates the suppressive effect of the compounds of this invention on internode elongation of aquatic rice plants.

5           3 hills (each hill comprising two seedlings) of aquatic rice seedlings ("Kinmaze" Variety) at the 3 leaf stage were transplanted in a 1/2000 are Wagner pot. 35 Days after transplanting, a dilute aqueous emulsion containing the test compound which had been prepared by diluting with  
10 water an emulsifiable concentrate formulated as in Example 17 was added dropwise into the flooding water in the pot at an application rate of 1 kg of the active ingredient per hectare.

3 Months after the application, assessments were made  
15 on the stem length, length of each of the first to fifth internodes and weight of each unit internode. The test results are indicated in Table 9 in terms of the percentages based on those in the untreated plots. The "unit internode weight" means dry weight of the internode of 1 cm long.

20

Table 9

Compound No. tested	Stem length	Each internode length					Unit internode weight				
		First	Second	Third	Fourth	Fifth	First	Second	Third	Fourth	Fifth
15	87	98	91	63	68	59	106	108	114	120	123

From the results of Table 9 as above, it will be seen that the new compounds of this invention exert the stunting effects on the elongation of the lower (third to fifth) internodes of aquatic rice plants and make these lower internodes thick and strong, so that the compounds are useful to reduce the risk of lodging of aquatic rice plants.

#### Example 25

This Example illustrates that the compounds of this invention reduce the risk of lodging of wheat plants.

10       Seeds of wheat ("Norin" No. 61 Variety) were sown in a field. When the wheat plants were grown to a stage of 30 to 40 cm in height, the field was divided into several plots each of 10 m<sup>2</sup> area. The wheat plants in each plot were treated by spraying their foliage evenly with a diluted aqueous preparations which had been prepared by formulating the compounds of this invention into a wettable powder as stated in Example 16 and diluting the wettable powder with such a volume of water that 500 litres of the diluted aqueous preparations so sprayed gave the rates of application of active ingredient as indicated in the following Table 10. For comparison purpose, 2-chloroethyl trimethyl ammonium chloride (CCC, known as chlormequat) was tested in the same manner as above.

20       80 Days after the treatment, the length of stem of the wheat plants and the extent of lodging were evaluated. The results obtained are tabulated in Table 10 below.

0123001

Table 10

	Compound No. tested	Rate of application of active ingredient (kg per hectare)	Stem length (%)	Extent of lodging
5	9	0.2	83	Very slight
		0.4	76	None
	15	0.2	80	Very slight
		0.4	75	None
	25	0.2	83	Very slight
10		0.4	77	None
	37	0.2	85	Slight
		0.4	80	Very slight
		0.5	101	None
	CCC (comparative)	1.0	81	Slight
		2.0	77	Very slight
	Untreated	-	100	Medium to great

15 Example 26

This Example demonstrates that the salts of the cyclohexane compounds of the formula (Ia) according to this invention have reduced influences on eyes in contact with the latter.

20 3 Rabbits (Female, "Japan White" Variety, 4 months-age) of which eyeballs were found to suffer from no damage by a check with 5% aqueous fluorescein sodium solution were fixed on a fixing table. 100 mg of the test compounds as such were

dropped into the left eyes of the rabbits and their eyelids were closed for one seconds, after which the rabbits were placed in a breeding case. The right eyes were remaining untreated for control. Eye-examinations were made 24 hours and 2, 3, 4, 7, 14 and 21 days after the treatment. The stimulus responses were evaluated in respect of various observations as indicated in Table 11, and the results are set out in Table 12.

Table 11

10 Cornea

(A) Opacity-Turbidity (the most opaque region of cornea was examined) were estimated by the following scales:

Scale 1 ... Sporadical or diffusive opacity occurred to such extent that iris still remained clearly visible.

15

Scale 2 ... Opacity occurred to such extent that fine parts of iris were somewhat dimly visible.

20

Scale 3 ... Opacity occurred to such extent that fine parts of iris were indistinct and the size of pupil is visually distinguishable with difficulty.

Scale 4 ... Opacity occurred to such extent that iris was not visible.

(B) Area suffering from corneal damage (denoted as "Area" below) was scaled as follows:

25

Scale 1 ...  $0 < \text{"Area"} < 1/4$





0123001

Scale 2 ...  $1/4 \leq \text{"Area"} < 1/2$

Scale 3 ...  $1/2 \leq \text{"Area"} < 3/4$

Scale 4 ...  $3/4 \leq \text{"Area"}$

A value of a product of scale (A) x scale (B) x 5  
5 was calculated and indicated as "Evaluation scores" of  
stimulus to cornea in the following Table 12, but this  
calculated value did not extend over the maximum value 80  
even for the worst case.

#### Iris

- 10 (A) Scale 1 ... one or more signs of acceleration of  
wrinkle formation, hyperemia, swelling  
and hyperemia of corneal surroundings in  
iris were detectable but reflection  
response of iris to light was still good.
- 15 Scale 2 ... one or more signs of no reflection  
response of iris to light, bleeding and  
wide-spread breakdown were detectable.

A value of a product of scale (A) x 5 was calculated  
and indicated as "Evaluation scores" of stimulus to iris in  
20 Table 12, but this calculated value did not extend over the  
maximum value 10 even for the worst case.

#### Conjunctiva

- (A) Redness (only the redness developed in the eyelid  
conjunctiva was examined.)
- 25 Scale 1 ... Acceleration of hyperemia was observed.

Scale 2 ... Wide-spread, deep-color redness and difficulty in identification of vasa were observed.

5 Scale 3 ... Wholly-spread, deep-color redness was observed.

(B) Chemosis

Scale 1 ... Acceleration of chemosis (including nictitating membrane) was observed.

10 Scale 2 ... Swelling associated with partial valgum of eyelid was observed.

Scale 3 ... Eyelid closing (in about half region) associated with swelling was observed.

Scale 4 ... Eyelid closing (about half to whole region) with swelling was observed.

15 (C) Secreta

Scale 1 ... More than ordinary amount (excluding the small quantity of secreta usually observed in inner canthus of healthy animal) of secreta was observed.

20 Scale 2 ... Secreta was formed in the amount to wet and stain eyelid and also hairs contact with eyelid.

Scale 3 ... Secreta was formed in the amount to wet and stain eyelid and also the eye surroundings over a fairly wide region.

25 A value of a product of [scale (A) + scale (B) + scale (C)] x 2 was calculated and indicated as "Evaluation

0123001

scores" in term of secreta in Table 12, but this calculated value did not extend over the maximum value 20 for the worst case.

Table 12

5	Compound No. tested	Compound No. 54			Compound No. 74			Compound No. 93			Compound No. 94			Compound No.9 (comparative)		
		2.5	2.6	2.5	2.7	3.0	2.7	3.0	2.6	2.5	2.5	2.5	2.1	2.7	2.6	2.6
10	Evaluation scores of stimulus	Cornea	Body weight(kg)													
			24 hours	0	0	0	0	5	0	0	0	0	0	0	15	20
			2 days	0	0	0	0	0	0	0	0	0	0	20	15	20
			3 days	0	0	0	0	0	0	0	0	0	0	20	10	15
			4 days	0	0	0	0	0	0	0	0	0	0	20	10	20
			7 days	0	0	0	0	0	0	0	0	0	0	20	5	5
			14 days	0	0	0	0	0	0	0	0	0	0	5	0	0
			21 days	0	0	0	0	0	0	0	0	0	0	0	0	0
			Average	0			0.2	0			0			11.4		
15	Evaluation scores of stimulus	Iris	24 hours	0	0	0	0	5	0	0	0	0	0	5	0	0
			2 days	0	0	0	0	0	0	0	0	0	0	5	0	0
			3 days	0	0	0	0	0	0	0	0	0	0	5	0	0
			4 days	0	0	0	0	0	0	0	0	0	0	0	0	0
			7 days	0	0	0	0	0	0	0	0	0	0	0	0	0
			14 days	0	0	0	0	0	0	0	0	0	0	0	0	0
			21 days	0	0	0	0	0	0	0	0	0	0	0	0	0
			Average	0			0.2	0			0			0.7		
25	Evaluation scores of stimulus	Conjunctiva	24 hours	4	2	2	2	4	2	6	4	6	0	8	8	6
			2 days	0	0	0	0	0	0	0	0	0	0	6	4	4
			3 days	0	0	0	0	0	0	0	0	0	0	2	0	0

to be cont'd...


0123001

5	Conjunctiva	4 days	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0
		7 days	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		14 days	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		21 days	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Average	0.4			0.4			0.8			0			1.9				

0123001

CLAIMS :

1. A cyclohexane compound of the general formula:




wherein R represents a hydrogen atom or an alkyl group, an alkylthioalkyl group or an unsubstituted or substituted phenyl group; and R<sup>1</sup> represents an alkyl group, an unsubstituted or substituted benzyl group, a phenethyl group, a phenoxyethyl group, a 2-thienylethyl group, an alkoxyethyl group or an alkylthioethyl group, or a salt of said cyclohexane compound.

2. A compound according to Claim 1 in which R is a hydrogen atom or an alkyl group.
3. A compound according to Claim 1 in which R<sup>1</sup> is an alkyl group.
4. A compound according to Claim 1 in which both R and R<sup>1</sup> are each an alkyl group of 1-8 carbon atoms.
5. A compound according to Claim 1 in which R is an alkyl group which is chosen from methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl and isopentyl.
6. A compound according to Claim 1 in which R<sup>1</sup> is an alkyl group which is chosen from methyl, ethyl, n-propyl, isopropyl.

7. A compound according to Claim 1 which is selected from 3,5-dioxo-4-propionylcyclohexanecarboxylic acid methyl ester; 3,5-dioxo-4-propionylcyclohexanecarboxylic acid ethyl ester; 3,5-dioxo-4-propionylcyclohexanecarboxylic acid n-propyl ester; 3,5-dioxo-4-butyrylcyclohexanecarboxylic acid; 3,5-dioxo-4-butyrylcyclohexanecarboxylic acid ethyl ester; 3,5-dioxo-4-butyrylcyclohexanecarboxylic acid n-propyl ester; 3,5-dioxo-4-butyrylcyclohexanecarboxylic acid isopropyl ester; 3,5-dioxo-4-butyrylcyclohexanecarboxylic acid n-butyl ester; 3,5-dioxo-4-butyrylcyclohexanecarboxylic acid isopentyl ester; 3,5-dioxo-4-isobutyrylcyclohexanecarboxylic acid ethyl ester; and 3,5-dioxo-4-acetylcyclohexanecarboxylic acid.


8. A compound as claimed in Claim 1, which is a salt of a cyclohexane compound of the general formula:



wherein  $R^2$  represents a hydrogen atom or an alkyl group and  $R^3$  represents an alkyl group.


0123001

9. A compound according to Claim 8 which is a salt of the cyclohexane compound represented by the general formula:




wherein  $R^2$  and  $R^3$  are as defined in Claim 8 and M represents an organic or inorganic cation.

10. A compound according to Claim 8 which is a salt of the cyclohexane compound represented by the general formula:



wherein  $R^3$  is as defined in Claim 8 and M represents an organic or inorganic cation.

11. A compound according to Claim 8 which is a salt of the cyclohexane compound represented by the general formula:



wherein  $R^3$  is as defined in Claim 8 and M represents an organic or inorganic cation.

12. A compound according to Claim 8 which is a salt of the cyclohexane compound of the formula (Ia) with an inorganic cation which is selected from cations of an alkali metal, an alkaline earth metal, aluminum, nickel, copper, manganese, cobalt, zinc, iron and silver, as well as ammonium cation.

13. A compound according to Claim 8 which is a salt of the cyclohexane compound of the formula (Ia) with an organic cation which is a substituted ammonium cation, the substituent on the ammonium cation being an alkyl, a hydroxy-alkyl, an alkenyl, benzyl, a halo-benzyl, pyridyl, an alkylpyridyl, or a polymethylene link optionally interrupted by an oxygen atom.

14. A compound according to Claim 8 which is chosen from:-


[bis-(2-hydroxyethyl)]ammonium salt of ethyl 3,5-dioxo-4-propionyl-cyclohexanecarboxylate; sodium salt of ethyl 3,5-dioxo-4-propionyl-cyclohexanecarboxylate; disodium salt of 3,5-dioxo-4-propionyl-cyclohexanecarboxylic acid; bis-  
-{[bis-(2-hydroxyethyl)]ammonium} salt of 3,5-dioxo-4-propionyl-cyclohexanecarboxylic acid; and [bis-(2-hydroxypropyl)]ammonium salt of ethyl 3,5-dioxo-4-propionyl-cyclohexanecarboxylate.






0123001

15. A process for the production of a cyclohexane compound as claimed in Claim 1 and represented by the general formula (I):




wherein R represents a hydrogen atom or an alkyl group, an alkylthioalkyl group or an unsubstituted or substituted phenyl group; and R<sup>1</sup> represents an alkyl group, an unsubstituted or substituted benzyl group, a phenethyl group, a phoxymethyl group, a 2-thienylmethyl group, an alkoxy-methyl group or an alkylthiomethyl group, or a salt of said cyclohexane compound, which comprises the steps of:-

(a) reacting a compound of the formula (III):



or a compound of the formula (IV):




wherein R is as defined above and M' is an alkali metal atom, with an organic acid chloride compound of the formula (V):

0123001




wherein  $\text{R}^1$  is as defined above, in an organic solvent and/or water as the reaction medium in the presence or absence of a base as an acid-binding agent to produce the compound of the formula (VI):



wherein R and  $\text{R}^1$  are as defined above,

(b) subjecting the compound of the formula (VI) to intermolecular rearrangement in the presence of a catalyst which is chosen from 4-N,N-dimethylaminopyridine, 4-N,N-diethylaminopyridine, 4-pyrrolidinoaminopyridine, N-methylimidazole and N-ethylimidazole to produce the compound of the formula (I):



wherein R and  $\text{R}^1$  are as defined above, and


(c) if desired, reacting the compound of the formula (I) with an inorganic or organic cation to produce the salt of the compound (I) with said inorganic or organic cation.

0123001

16. A process according to Claim 15, in which the step of reacting the compound (III) or (IV) with an organic acid chloride (V) to produce the compound (VI) is carried out in a mixture of water and an organic solvent and in the presence of  $\gamma$ -picoline as a phase transition catalyst.


17. A process according to Claim 15, in which the step of subjecting the compound (VI) to rearrangement to produce the compound (I) is carried out in the presence of 4-N,N-dimethylaminopyridine as the catalyst.

18. A plant-growth regulating composition comprising as active ingredient a cyclohexane compound of the general formula:




wherein R represents a hydrogen atom or an alkyl group, an alkylthioalkyl group or an unsubstituted or substituted phenyl group; and R<sup>1</sup> represents an alkyl group, an unsubstituted or substituted benzyl group, a phenethyl group, a phoxymethyl group, a 2-thienylmethyl group, an alkoxy-methyl group or an alkylthiomethyl group, or a salt of said cyclohexane compound.

19. A plant-growth regulating composition comprising as active ingredient a salt of a cyclohexane compound having the general formula:




wherein  $R^2$  represents a hydrogen atom or an alkyl group and  $R^3$  represents an alkyl group.

20. A composition according to Claim 19, in which the salt is of the general formula:



wherein  $R^2$  and  $R^3$  are as defined in Claim 19 and M represents an organic or inorganic cation.


21. A composition according to Claim 19, in which the salt is of the formula:



wherein  $R^3$  and M are as defined in Claim 20.

0123001

22. A composition according to Claim 19, in which the salt is of the formula:



wherein R<sup>3</sup> and M are as defined in Claim 20.

23. A method of regulating the growth of plant, which comprises applying an effective amount of a cyclohexane compound of the formula (I) or a salt thereof as claimed in Claim 1 to the foliage or seed of the plant to be treated or to the soil or locus where the plant to be treated is grown.

FIG. I




FIG.2




FIG. 3






FIG.4




FIG. 5




FIG. 6





FIG. 7





European Patent  
Office

# EUROPEAN SEARCH REPORT

0123001

Application number

EP 83 40 1861

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 3)
A	US-A-4 281 204 (B.J. WILLIS) * Column 3, line 43 - column 5, line 55 *	1	C 07 C 69/757 C 07 C 62/20 C 07 C 149/30 C 07 C 149/28 C 07 D 333/24
A	--- CHEMICAL ABSTRACTS, vol. 88, 1978, page 466, no. 74127k, Columbus, Ohio, USA; & JP - A - 77 118 448 (NIPPON SODA CO., LTD.) 04-10-1977 * Abstract *	1	C 07 C 91/26 C 07 C 87/30 C 07 C 67/343 A 01 N 37/10
A	--- US-A-3 950 420 (MIKIO SAWAKI) * Column 3, lines 30-58 *	1	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 3)
			C 07 C 69/00 C 07 C 67/00 C 07 C 62/00
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 26-05-1984	Examiner KINZINGER J.M.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone		T : theory or principle underlying the invention	
Y : particularly relevant if combined with another document of the same category		E : earlier patent document, but published on, or after the filing date	
A : technological background		D : document cited in the application	
O : non-written disclosure		L : document cited for other reasons	
P : intermediate document		& : member of the same patent family, corresponding document	

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☒ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**